arXiv:0810.0058v1 [nlin.SI] 1 Oct 2008

Lax forms of the q-Painlevé equations

Mikio Murata
Department of Physics and Mathematics,
Aoyama Gakuin University,
5-10-1 Fuchinobe Sagamihara-shi,
Kanagawa 229-8558, Japan

Abstract

All q-Painlevé equations which are obtained from the q-analog of the sixth Painlevé equation are expressed in a Lax formalism. They are characterized by the data of the associated linear q-difference equations. The degeneration pattern from the q-Painlevé equation of type A_2 is also presented.

Keywords: Painlevé equations, q-difference equations, Completely integrable systems.

2000 Mathematics Subject Classification: 33E17, 34M55, 39A12.

1 Introduction

Discrete Painlevé equations are studied from various points of view as integrable systems ([5]). They are discrete equations which are reduced to the Painlevé equations in suitable limiting process, and moreover, which pass the singularity confinement test. Passing this test can be thought of as a difference analogue of the Painlevé property. The singularity confinement test has been proposed by Grammaticos et al. as a criterion for the integrability of discrete dynamical systems ([2]).

Discrete Painlevé equations were classified on the basis of the types of rational surfaces connected to extended affine Weyl groups ([9, 10]). There are three types of discrete Painlevé equations: elliptic-difference, q-difference and difference. We especially give the list of q-Painlevé equations in the discrete Painlevé equations. See Table 1. As is well-known, the sixth Painlevé equation yields the other five Painlevé equations by a process of coalescence.

Abbrev.	q-$P(A_0^*)$	q-$P(A_1)$	q-$P(A_2)$	q-$P(A_3)$	q-$P(A_4)$	q-$P(A_5)$	q-$P(A_5)^\sharp$	q-$P(A_6)$	q-$P(A_6)^\sharp$	q-$P(A_7)$	q-$P(A_7')$
Surface	$A_0^{(1)*}$	$A_1^{(1)}$	$A_2^{(1)}$	$A_3^{(1)}$	$A_4^{(1)}$	$A_5^{(1)}$	$A_5^{(1)}$	$A_6^{(1)}$	$A_6^{(1)}$	$A_7^{(1)}$	$A_7^{(1)\prime}$
Symmetry	$E_8^{(1)}$	$E_7^{(1)}$	$E_6^{(1)}$	$D_5^{(1)}$	$A_4^{(1)}$	$(A_2+A_1)^{(1)}$	$(A_2+A_1)^{(1)}$	$(A_1+A_1)^{(1)}$	$(A_1+A_1)^{(1)}$	$A_1^{(1)}$	$A_1^{(1)}$

Table 1: The q-Painlevé equations

Among the q-Painlevé equations, the q-Painlevé equation of type A_0^* (q-$P(A_0^*)$) is the most generic one because the other q-Painlevé equations can be obtained from this equation by limiting procedure. These equations are organized in a degeneration pattern obtained through coalescence. See Table 2.

$$\begin{tikzcd}
q\text{-}P(A_0^*) \arrow[r] & q\text{-}P(A_1) \arrow[r] & q\text{-}P(A_2) \arrow[r] & q\text{-}P(A_3) \arrow[r] & {} \\
{} \arrow[r] & q\text{-}P(A_4) \arrow[r] \arrow[dr] & q\text{-}P(A_5) \arrow[r] & q\text{-}P(A_6) \arrow[r] & q\text{-}P(A_7) \\
& & q\text{-}P(A_5)^\sharp \arrow[r] \arrow[ur] & q\text{-}P(A_6)^\sharp \arrow[r] \arrow[ur] & q\text{-}P(A_7')
\end{tikzcd}$$

Table 2: Degeneration pattern for the q-Painlevé equations

Another important aspect of the Painlevé equations is their connection to the monodromy-preserving deformation of linear differential equations. The generalized Riemann problem was already studied for linear differential, difference and q-difference equations in the Birkhoff's paper, [1]. Jimbo and Sakai studied the deformation of a 2×2 matrix system of q-difference equations and found the q-Painlevé equation of type A_3 (q-$P(A_3)$), which is commonly known as q-$\mathrm{P_{VI}}$ ([4]). Sakai also found a Lax form of the q-Painlevé equation of type A_2 (q-$P(A_2)$), a particular case of a q-Garnier system ([7, 8]). Hay et al. found Lax forms of q-Painlevé equations, reductions from a Lax pair for a lattice modified KdV equation ([3]). However, Lax forms of a lot of q-Painlevé equations have not been obtained yet.

In this paper, we present Lax pairs of all q-Painlevé equations which are obtained from q-$P(A_3)$. In Section 2, we illustrate the connection preserving deformation and derive q-$P(A_3)$. We also propose Lax pairs of q-$P(A_4)$, q-$P(A_5)$, q-$P(A_5)^\sharp$, q-$P(A_6)$, q-$P(A_6)^\sharp$, q-$P(A_7)$ and q-$P(A_7')$. In Section 3, we give replacements of the parameters for the degeneration. The Lax form of q-$P(A_3)$ can be obtained from the Lax form of q-$P(A_2)$. In Section 4, we give the Lax form of q-$P(A_2)$ and replacements of the parameters for the degeneration.

2 Lax forms of q-Painlevé equations

2.1 Derivation of q-$P(A_3)$

In this section, we illustrate the connection preserving deformation and derive the q-Painlevé equation of type A_3.

Consider a 2×2 matrix system with polynomial coefficients

$$Y(qx,t) = A(x,t)Y(x,t). \tag{2.1}$$

The connection preserving deformation of the linear q-difference equation, which is a discrete counterpart of monodromy preserving deformation, is equivalent to existence of a linear deformation equation whose coefficients are rational in x. We express the deformation equation in the form

$$Y(x,qt) = B(x,t)Y(x,t). \tag{2.2}$$

The compatibility condition for the systems (2.1) and (2.2) reads

$$A(x,qt)B(x,t) = B(qx,t)A(x,t). \tag{2.3}$$

q-$P(A_3)$ can be obtained from the condition (2.3).

We take $A(x,t)$ to be of the form

$$A(x,t) = A_0(t) + xA_1(t) + x^2A_2, \tag{2.4}$$

$$A_2 = \begin{pmatrix} \kappa_1 & 0 \\ 0 & \kappa_2 \end{pmatrix}, \quad A_0(t) \text{ has eigenvalues } \theta_1 t,\ \theta_2 t, \tag{2.5}$$

$$\det A(x,t) = \kappa_1\kappa_2(x-a_1t)(x-a_2t)(x-a_3)(x-a_4). \tag{2.6}$$

Here the parameters κ_j, θ_j, a_j are independent of t. We have

$$\kappa_1\kappa_2a_1a_2a_3a_4 = \theta_1\theta_2. \tag{2.7}$$

Define $y = y(t)$, $z_i = z_i(t)$ $(i = 1, 2)$ by

$$A_{12}(y,t) = 0, \quad A_{11}(y,t) = \kappa_1z_1, \quad A_{22}(y,t) = \kappa_2z_2, \tag{2.8}$$

so that

$$z_1z_2 = \kappa_1\kappa_2(y-a_1t)(y-a_2t)(y-a_3)(y-a_4). \tag{2.9}$$

The matrix $A(x,t)$ can be parametrized as

$$A(x,t)=\begin{pmatrix}\kappa_1((x-y)(x-\alpha)+z_1) & \kappa_2 w(x-y)\\ \kappa_1 w^{-1}(\gamma x+\delta) & \kappa_2((x-y)(x-\beta)+z_2)\end{pmatrix}. \tag{2.10}$$

Here

$$\begin{aligned}\alpha=&\frac{1}{\kappa_1-\kappa_2}[y^{-1}((\theta_1+\theta_2)t-\kappa_1 z_1-\kappa_2 z_2)\\ &-\kappa_2((a_1+a_2)t+a_3+a_4-2y)],\end{aligned} \tag{2.11}$$

$$\begin{aligned}\beta=&\frac{1}{\kappa_1-\kappa_2}[-y^{-1}((\theta_1+\theta_2)t-\kappa_1 z_1-\kappa_2 z_2)\\ &+\kappa_1((a_1+a_2)t+a_3+a_4-2y)],\end{aligned} \tag{2.12}$$

$$\begin{aligned}\gamma=&z_1+z_2+(y+\alpha)(y+\beta)+(\alpha+\beta)y-a_1a_2t^2\\ &-(a_1+a_2)(a_3+a_4)t-a_3a_4,\end{aligned} \tag{2.13}$$

$$\delta=y^{-1}(a_1a_2a_3a_4t^2-(\alpha y+z_1)(\beta y+z_2)). \tag{2.14}$$

The quantity $w=w(t)$ is related to the ‘gauge’ freedom, and does not enter the final result for the q-$P(A_3)$. The matrix $B(x,t)$ is a rational function of the form

$$B(x,t)=\frac{x}{(x-a_1qt)(x-a_2qt)}(xI+B_0(t)). \tag{2.15}$$

The compatibility (2.3) is equivalent to

$$A(a_iqt,qt)(a_iqtI+B_0(t))=0 \quad (i=1,2), \tag{2.16}$$

$$(a_iqtI+B_0(t))A(a_it,t)=0 \quad (i=1,2), \tag{2.17}$$

$$A_0(qt)B_0(t)=qB_0(t)A_0(t). \tag{2.18}$$

Substituting the parametrization above, one obtains a set of q-difference equations. Let us use the notations $\bar{y}=y(qt)$ and so forth. Introduce z by

$$z_1=\frac{(y-a_1t)(y-a_2t)}{\kappa_1qz}, \quad z_2=\kappa_1qz(y-a_3)(y-a_4). \tag{2.19}$$

Then the matrix $B_0(t) = (B_{ij})$ is parametrized as follows:

$$\begin{aligned}
B_{11} &= \frac{-\kappa_2 q\bar{z}}{1-\kappa_2\bar{z}}\left(-\beta + \frac{t(a_1+a_2)-y}{\kappa_2\bar{z}}\right),\\
B_{12} &= \frac{\kappa_2 q w\bar{z}}{1-\kappa_2\bar{z}},\\
B_{21} &= \frac{\kappa_1 q\bar{z}}{w(1-\kappa_1 q\bar{z})}\left(a_1qt-\bar{\alpha}+\frac{a_2qt-\bar{y}}{\kappa_1 q\bar{z}}\right)\left(a_1t-\beta+\frac{a_2t-y}{\kappa_2\bar{z}}\right)\\
&= \frac{\kappa_1 q\bar{z}}{w(1-\kappa_1 q\bar{z})}\left(a_2qt-\bar{\alpha}+\frac{a_1qt-\bar{y}}{\kappa_1 q\bar{z}}\right)\left(a_2t-\beta+\frac{a_1t-y}{\kappa_2\bar{z}}\right),\\
B_{22} &= \frac{-\kappa_1 q\bar{z}}{1-\kappa_1 q\bar{z}}\left(-\bar{\alpha} + \frac{qt(a_1+a_2)-\bar{y}}{\kappa_1 q\bar{z}}\right).
\end{aligned}$$

Set further

$$b_1 = \frac{a_1a_2}{\theta_1}, \quad b_2 = \frac{a_1a_2}{\theta_2}, \quad b_3 = \frac{1}{\kappa_1 q}, \quad b_4 = \frac{1}{\kappa_2}. \tag{2.20}$$

Equations (2.16)–(2.18) are equivalent to

$$\frac{y\bar{y}}{a_3a_4} = \frac{(\bar{z}-b_1t)(\bar{z}-b_2t)}{(\bar{z}-b_3)(\bar{z}-b_4)}, \tag{2.21}$$

$$\frac{z\bar{z}}{b_3b_4} = \frac{(y-a_1t)(y-a_2t)}{(y-a_3)(y-a_4)}, \tag{2.22}$$

$$\frac{\bar{w}}{w} = \frac{b_4}{b_3}\frac{\bar{z}-b_3}{\bar{z}-b_4}. \tag{2.23}$$

We have a single constraint

$$\frac{b_1b_2}{b_3b_4} = q\frac{a_1a_2}{a_3a_4}. \tag{2.24}$$

q-$P(A_3)$ is (2.21) and (2.22).

2.2 Lax form of q-$P(A_4)$

Consider a 2×2 matrix system with polynomial coefficients

$$Y(qx,t) = A(x,t)Y(x,t). \tag{2.25}$$

We express the deformation equation in the form

$$Y(x,qt) = B(x,t)Y(x,t) \tag{2.26}$$

and can express the q-Painlevé equation of type A_4 in the form

$$A(x, qt)B(x, t) = B(qx, t)A(x, t) \tag{2.27}$$

by the compatibility of the deformation equation and the original linear q-difference equation.

We take $A(x, t)$ to be of the form

$$A(x, t) = A_0(t) + xA_1(t) + x^2 A_2, \tag{2.28}$$

$$A_2 = \begin{pmatrix} \kappa_1 & 0 \\ 0 & 0 \end{pmatrix}, \quad A_0(t) \text{ has eigenvalues } \theta_1 t,\ \theta_2 t, \tag{2.29}$$

$$\det A(x, t) = \kappa_1\kappa_2(x - a_1 t)(x - a_2 t)(x - a_3). \tag{2.30}$$

We have

$$-\kappa_1\kappa_2 a_1 a_2 a_3 = \theta_1\theta_2. \tag{2.31}$$

The matrix $B(x, t)$ is a rational function of the form

$$B(x, t) = \frac{x}{(x - a_1 qt)(x - a_2 qt)}(xI + B_0(t)). \tag{2.32}$$

Define $y = y(t)$, $z_i = z_i(t)$ $(i = 1, 2)$ by

$$A_{12}(y, t) = 0, \quad A_{11}(y, t) = \kappa_1 z_1, \quad A_{22}(y, t) = z_2, \tag{2.33}$$

so that

$$z_1 z_2 = \kappa_2(y - a_1 t)(y - a_2 t)(y - a_3). \tag{2.34}$$

The matrix $A(x, t)$ can be parametrized as

$$A(x, t) = \begin{pmatrix} \kappa_1((x - y)(x - \alpha) + z_1) & w(x - y) \\ \kappa_1 w^{-1}(\gamma x + \delta) & \kappa_2(x - y) + z_2 \end{pmatrix}. \tag{2.35}$$

Here

$$\alpha = \frac{1}{\kappa_1}[y^{-1}((\theta_1 + \theta_2)t - \kappa_1 z_1 - z_2) + \kappa_2], \tag{2.36}$$

$$\gamma = z_2 - \kappa_2((2y + \alpha) - (a_1 + a_2)t - a_3), \tag{2.37}$$

$$\delta = y^{-1}(-\kappa_2 a_1 a_2 a_3 t^2 - (\alpha y + z_1)(-\kappa_2 y + z_2)). \tag{2.38}$$

Introduce z by

$$z_1 = \frac{(y - a_1 t)(y - a_2 t)}{\kappa_1 qz}, \quad z_2 = \kappa_1\kappa_2 qz(y - a_3). \tag{2.39}$$

Then the matrix $B_0(t) = (B_{ij})$ is parametrized as follows:

$$B_{11} = -q\bar{z}\left(\kappa_2 + \frac{t(a_1+a_2)-y}{\bar{z}}\right), \tag{2.40}$$

$$B_{12} = qw\bar{z}, \tag{2.41}$$

$$\begin{aligned} B_{21} &= \frac{\kappa_1 q\bar{z}}{w(1-\kappa_1 q\bar{z})}\left(a_1qt - \bar{\alpha} + \frac{a_2qt-\bar{y}}{\kappa_1 q\bar{z}}\right)\left(\kappa_2 + \frac{a_2t-y}{\bar{z}}\right) \\ &= \frac{\kappa_1 q\bar{z}}{w(1-\kappa_1 q\bar{z})}\left(a_2qt - \bar{\alpha} + \frac{a_1qt-\bar{y}}{\kappa_1 q\bar{z}}\right)\left(\kappa_2 + \frac{a_1t-y}{\bar{z}}\right), \end{aligned} \tag{2.42}$$

$$B_{22} = \frac{-\kappa_1 q\bar{z}}{1-\kappa_1 q\bar{z}}\left(-\bar{\alpha} + \frac{qt(a_1+a_2)-\bar{y}}{\kappa_1 q\bar{z}}\right). \tag{2.43}$$

Set further

$$b_1 = \frac{a_1a_2}{\theta_1}, \quad b_2 = \frac{a_1a_2}{\theta_2}, \quad b_3 = \frac{1}{\kappa_1 q}, \quad a_4 = -\kappa_2. \tag{2.44}$$

Equation (2.27) are equivalent to

$$\frac{y\bar{y}}{a_3a_4} = -\frac{(\bar{z}-b_1t)(\bar{z}-b_2t)}{\bar{z}-b_3}, \tag{2.45}$$

$$\frac{z\bar{z}}{b_3} = -\frac{(y-a_1t)(y-a_2t)}{a_4(y-a_3)}, \tag{2.46}$$

$$\frac{\bar{w}}{w} = -\frac{\bar{z}}{b_3} + 1. \tag{2.47}$$

We have a constraint

$$\frac{b_1b_2}{b_3} = q\frac{a_1a_2}{a_3a_4}. \tag{2.48}$$

q-$P(A_4)$ is (2.45) and (2.46).

2.3 Lax form of q-$P(A_5)$

Consider a 2×2 matrix system with polynomial coefficients

$$Y(qx,t) = A(x,t)Y(x,t). \tag{2.49}$$

We express the deformation equation

$$Y(x,qt) = B(x,t)Y(x,t) \tag{2.50}$$

and can express the q-Painlevé equation of type A_5 in the form

$$A(x,qt)B(x,t) = B(qx,t)A(x,t) \tag{2.51}$$

by the compatibility of the deformation equation and the original linear q-difference equation.

We take $A(x,t)$ to be of the form

$$A(x,t) = A_0(t) + xA_1(t) + x^2 A_2, \tag{2.52}$$

$$A_2 = \begin{pmatrix} \kappa_1 & 0 \\ 0 & 0 \end{pmatrix}, \quad A_0(t) \text{ has eigenvalues } \theta_1 t,\ 0, \tag{2.53}$$

$$\det A(x,t) = \kappa_1\kappa_2 x(x-a_1t)(x-a_2t). \tag{2.54}$$

The matrix $B(x,t)$ is a rational function of the form

$$B(x,t) = \frac{x}{(x-a_1qt)(x-a_2qt)}(xI + B_0(t)). \tag{2.55}$$

Define $y = y(t)$, $z_i = z_i(t)$ $(i = 1,2)$ by

$$A_{12}(y,t) = 0, \quad A_{11}(y,t) = \kappa_1 z_1, \quad A_{22}(y,t) = z_2, \tag{2.56}$$

so that

$$z_1 z_2 = \kappa_2 y(y-a_1t)(y-a_2t). \tag{2.57}$$

The matrix $A(x,t)$ can be parametrized as

$$A(x,t) = \begin{pmatrix} \kappa_1((x-y)(x-\alpha)+z_1) & w(x-y) \\ \kappa_1 w^{-1}(\gamma x+\delta) & \kappa_2(x-y)+z_2 \end{pmatrix}. \tag{2.58}$$

Here

$$\alpha = \frac{1}{\kappa_1}[y^{-1}(\theta_1 t - \kappa_1 z_1 - z_2) + \kappa_2], \tag{2.59}$$

$$\gamma = z_2 - \kappa_2(2y + \alpha - t(a_1+a_2)), \tag{2.60}$$

$$\delta = -y^{-1}(\alpha y + z_1)(-\kappa_2 y + z_2). \tag{2.61}$$

Introduce z by

$$z_1 = \frac{(y-a_1t)(y-a_2t)}{\kappa_1 qz}, \quad z_2 = \kappa_1\kappa_2 qzy. \tag{2.62}$$

Then the matrix $B_0(t) = (B_{ij})$ is parametrized as follows:

$$B_{11} = -q\bar{z}\left(\kappa_2 + \frac{t(a_1+a_2) - y}{\bar{z}}\right), \tag{2.63}$$

$$B_{12} = qw\bar{z}, \tag{2.64}$$

$$\begin{aligned} B_{21} &= \frac{\kappa_1 q\bar{z}}{w(1-\kappa_1 q\bar{z})}\left(a_1qt - \bar{\alpha} + \frac{a_2qt - \bar{y}}{\kappa_1 q\bar{z}}\right)\left(\kappa_2 + \frac{a_2t - y}{\bar{z}}\right) \\ &= \frac{\kappa_1 q\bar{z}}{w(1-\kappa_1 q\bar{z})}\left(a_2qt - \bar{\alpha} + \frac{a_1qt - \bar{y}}{\kappa_1 q\bar{z}}\right)\left(\kappa_2 + \frac{a_1t - y}{\bar{z}}\right), \end{aligned} \tag{2.65}$$

$$B_{22} = \frac{-\kappa_1 q\bar{z}}{1-\kappa_1 q\bar{z}}\left(-\bar{\alpha} + \frac{qt(a_1+a_2) - \bar{y}}{\kappa_1 q\bar{z}}\right). \tag{2.66}$$

Set further

$$b_1 = \frac{a_1a_2}{\theta_1}, \quad b_2 = -\frac{\theta_1}{\kappa_1\kappa_2}, \quad b_3 = \frac{1}{\kappa_1 q}, \quad a_4 = -\kappa_2. \tag{2.67}$$

Equation (2.51) are equivalent to

$$\frac{y\bar{y}}{a_3a_4} = \frac{b_2t(\bar{z} - b_1t)}{\bar{z} - b_3}, \tag{2.68}$$

$$\frac{z\bar{z}}{b_3} = -\frac{(y-a_1t)(y-a_2t)}{a_4y}, \tag{2.69}$$

$$\frac{\bar{w}}{w} = -\frac{\bar{z}}{b_3} + 1. \tag{2.70}$$

We have a constraint

$$\frac{b_1b_2}{b_3} = q\frac{a_1a_2}{a_4}. \tag{2.71}$$

q-$P(A_5)$ is (2.68) and (2.69).

2.4 Lax form of q-$P(A_5)^\sharp$

Consider a 2×2 matrix system with polynomial coefficients

$$Y(qx,t) = A(x,t)Y(x,t). \tag{2.72}$$

We express the deformation equation

$$Y(x,qt) = B(x,t)Y(x,t) \tag{2.73}$$

and can express the q-Painlevé equation of type ${A_5}^\sharp$ in the form

$$A(x,qt)B(x,t) = B(qx,t)A(x,t) \tag{2.74}$$

by the compatibility of the deformation equation and the original linear q-difference equation.

We take $A(x,t)$ to be of the form

$$A(x,t) = A_0(t) + xA_1(t) + x^2A_2, \tag{2.75}$$

$$A_2 = \begin{pmatrix} \kappa_1 & 0 \\ 0 & 0 \end{pmatrix}, \quad A_0(t) \text{ has eigenvalues } \theta_1 t,\ 0, \tag{2.76}$$

$$\det A(x,t) = \kappa_1\kappa_2 x(x-a_1t)(x-a_3). \tag{2.77}$$

The matrix $B(x,t)$ is a rational function of the form

$$B(x,t) = \frac{1}{x-a_1qt}(xI + B_0(t)). \tag{2.78}$$

Define $y = y(t)$, $z_i = z_i(t)$ $(i = 1,2)$ by

$$A_{12}(y,t) = 0, \quad A_{11}(y,t) = \kappa_1 z_1, \quad A_{22}(y,t) = z_2, \tag{2.79}$$

so that

$$z_1z_2 = \kappa_2 y(y-a_1t)(y-a_3). \tag{2.80}$$

The matrix $A(x,t)$ can be parametrized as

$$A(x,t) = \begin{pmatrix} \kappa_1((x-y)(x-\alpha)+z_1) & w(x-y) \\ \kappa_1 w^{-1}(\gamma x+\delta) & \kappa_2(x-y)+z_2 \end{pmatrix}. \tag{2.81}$$

Here

$$\alpha = \frac{1}{\kappa_1}[y^{-1}(\theta_1 t - \kappa_1 z_1 - z_2) + \kappa_2], \tag{2.82}$$

$$\gamma = z_2 - \kappa_2(2y + \alpha - a_1t - a_3), \tag{2.83}$$

$$\delta = -y^{-1}(\alpha y + z_1)(-\kappa_2 y + z_2). \tag{2.84}$$

Introduce z by

$$z_1 = \frac{y(y-a_1t)}{\kappa_1 qz}, \quad z_2 = \kappa_1\kappa_2 qz(y-a_3). \tag{2.85}$$

Then the matrix $B_0(t) = (B_{ij})$ is parametrized as follows:

$$B_{11} = -q\bar{z}\left(\kappa_2 + \frac{a_1 t - y}{\bar{z}}\right), \tag{2.86}$$

$$B_{12} = qw\bar{z}, \tag{2.87}$$

$$\begin{aligned} B_{21} &= \frac{\kappa_1 q\bar{z}}{w(1-\kappa_1 q\bar{z})}\left(a_1 qt - \bar{\alpha} - \frac{\bar{y}}{\kappa_1 q\bar{z}}\right)\left(\kappa_2 - \frac{y}{\bar{z}}\right) \\ &= \frac{\kappa_1 q\bar{z}}{w(1-\kappa_1 q\bar{z})}\left(-\bar{\alpha} + \frac{a_1 qt - \bar{y}}{\kappa_1 q\bar{z}}\right)\left(\kappa_2 + \frac{a_1 t - y}{\bar{z}}\right), \end{aligned} \tag{2.88}$$

$$B_{22} = \frac{-\kappa_1 q\bar{z}}{1-\kappa_1 q\bar{z}}\left(-\bar{\alpha} + \frac{a_1 qt - \bar{y}}{\kappa_1 q\bar{z}}\right). \tag{2.89}$$

Set further

$$b_1 = \frac{a_1}{\theta_1}, \quad b_2 = -\frac{\theta_1}{\kappa_1\kappa_2 a_3}, \quad b_3 = \frac{1}{\kappa_1 q}, \quad a_4 = -\kappa_2. \tag{2.90}$$

Equation (2.74) are equivalent to

$$\frac{y\bar{y}}{a_3 a_4} = -\frac{\bar{z}(\bar{z} - b_2 t)}{\bar{z} - b_3}, \tag{2.91}$$

$$\frac{z\bar{z}}{b_3} = -\frac{y(y - a_1 t)}{a_4(y - a_3)}, \tag{2.92}$$

$$\frac{\bar{w}}{w} = -\frac{\bar{z}}{b_3} + 1. \tag{2.93}$$

We have a constraint

$$\frac{b_1 b_2}{b_3} = q\frac{a_1}{a_3 a_4}. \tag{2.94}$$

q-$P(A_5)^\sharp$ is (2.91) and (2.92).

2.5 Lax form of q-$P(A_6)$

Consider a 2×2 matrix system with polynomial coefficients

$$Y(qx, t) = A(x, t)Y(x, t). \tag{2.95}$$

We express the deformation equation

$$Y(x, qt) = B(x, t)Y(x, t) \tag{2.96}$$

and can express the q-Painlevé equation type A_6 in the form

$$A(x, qt)B(x, t) = B(qx, t)A(x, t) \tag{2.97}$$

by the compatibility of the deformation equation and the original linear q-difference equation.

We take $A(x,t)$ to be of the form

$$A(x,t) = A_0(t) + xA_1(t) + x^2A_2, \tag{2.98}$$

$$A_2 = \begin{pmatrix} \kappa_1 & 0 \\ 0 & 0 \end{pmatrix}, \quad A_0(t) \text{ has eigenvalues } \theta_1 t,\ 0, \tag{2.99}$$

$$\det A(x,t) = \kappa_1\kappa_2 x^2(x - a_1 t). \tag{2.100}$$

The matrix $B(x,t)$ is a rational function of the form

$$B(x,t) = \frac{1}{x - a_1 q t}(xI + B_0(t)). \tag{2.101}$$

Define $y = y(t)$, $z_i = z_i(t)$ $(i = 1,2)$ by

$$A_{12}(y,t) = 0, \quad A_{11}(y,t) = \kappa_1 z_1, \quad A_{22}(y,t) = z_2, \tag{2.102}$$

so that

$$z_1 z_2 = \kappa_2 y^2(y - a_1 t). \tag{2.103}$$

The matrix $A(x,t)$ can be parametrized as

$$A(x,t) = \begin{pmatrix} \kappa_1((x-y)(x-\alpha) + z_1) & w(x-y) \\ \kappa_1 w^{-1}(\gamma x + \delta) & \kappa_2(x-y) + z_2 \end{pmatrix}. \tag{2.104}$$

Here

$$\alpha = \frac{1}{\kappa_1}[y^{-1}(\theta_1 t - \kappa_1 z_1 - z_2) + \kappa_2], \tag{2.105}$$

$$\gamma = z_2 - \kappa_2(2y + \alpha - a_1 t), \tag{2.106}$$

$$\delta = -y^{-1}(\alpha y + z_1)(-\kappa_2 y + z_2). \tag{2.107}$$

Introduce z by

$$z_1 = \frac{y(y - a_1 t)}{\kappa_1 q z}, \quad z_2 = \kappa_1\kappa_2 q z y. \tag{2.108}$$

Then the matrix $B_0(t) = (B_{ij})$ is parametrized as follows:

$$B_{11} = -q\bar{z}\left(\kappa_2 + \frac{a_1 t - y}{\bar{z}}\right), \tag{2.109}$$

$$B_{12} = qw\bar{z}, \tag{2.110}$$

$$\begin{aligned} B_{21} &= \frac{\kappa_1 q\bar{z}}{w(1 - \kappa_1 q\bar{z})}\left(a_1 q t - \bar{\alpha} - \frac{\bar{y}}{\kappa_1 q\bar{z}}\right)\left(\kappa_2 - \frac{y}{\bar{z}}\right) \\ &= \frac{\kappa_1 q\bar{z}}{w(1 - \kappa_1 q\bar{z})}\left(-\bar{\alpha} + \frac{a_1 q t - \bar{y}}{\kappa_1 q\bar{z}}\right)\left(\kappa_2 + \frac{a_1 t - y}{\bar{z}}\right). \end{aligned} \tag{2.111}$$

$$B_{22} = \frac{-\kappa_1 q\bar{z}}{1 - \kappa_1 q\bar{z}}\left(-\bar{\alpha} + \frac{a_1 q t - \bar{y}}{\kappa_1 q\bar{z}}\right). \tag{2.112}$$

Set further

$$b_1 = \frac{a_1}{\theta_1}, \quad b_2 = -\frac{\theta_1}{\kappa_1 \kappa_2}, \quad b_3 = \frac{1}{\kappa_1 q}, \quad b_4 = -\kappa_2. \tag{2.113}$$

Equation (2.97) are equivalent to

$$\frac{y\bar{y}}{a_4} = \frac{b_2 t \bar{z}}{\bar{z} - b_3}, \tag{2.114}$$

$$\frac{z\bar{z}}{b_3} = -\frac{y(y - a_1 t)}{a_4 y}, \tag{2.115}$$

$$\frac{\bar{w}}{w} = -\frac{\bar{z}}{b_3} + 1. \tag{2.116}$$

We have a constraint

$$\frac{b_1 b_2}{b_3} = q\frac{a_1}{a_4}. \tag{2.117}$$

q-$P(A_6)$ is (2.114) and (2.115).

2.6 Lax form of q-$P(A_6)^\sharp$

Consider a 2×2 matrix system with polynomial coefficients

$$Y(qx, t) = A(x, t)Y(x, t). \tag{2.118}$$

We express the deformation equation

$$Y(x, qt) = B(x, t)Y(x, t) \tag{2.119}$$

and can express the q-Painlevé equation of type ${A_6}^\sharp$ in the form

$$A(x, qt)B(x, t) = B(qx, t)A(x, t) \tag{2.120}$$

by the compatibility of the deformation equation and the original linear q-difference equation.

We take $A(x, t)$ to be of the form

$$A(x, t) = A_0(t) + xA_1(t) + x^2 A_2, \tag{2.121}$$

$$A_2 = \begin{pmatrix} \kappa_1 & 0 \\ 0 & 0 \end{pmatrix}, \quad A_0(t) \text{ has eigenvalues } \theta_1 t,\ 0, \tag{2.122}$$

$$\det A(x, t) = \kappa_1 \kappa_2 x^2 (x - a_3), \tag{2.123}$$

The matrix $B(x, t)$ is a rational function of the form

$$B(x, t) = \frac{1}{x}(xI + B_0(t)). \tag{2.124}$$

Define $y = y(t)$, $z_i = z_i(t)$ $(i = 1, 2)$ by

$$A_{12}(y,t) = 0, \quad A_{11}(y,t) = \kappa_1 z_1, \quad A_{22}(y,t) = z_2, \tag{2.125}$$

so that

$$z_1 z_2 = \kappa_2 y^2 (y - a_3). \tag{2.126}$$

The matrix $A(x,t)$ can be parametrized as

$$A(x,t) = \begin{pmatrix} \kappa_1((x-y)(x-\alpha)+z_1) & w(x-y) \\ \kappa_1 w^{-1}(\gamma x + \delta) & \kappa_2(x-y)+z_2 \end{pmatrix}. \tag{2.127}$$

Here

$$\alpha = \frac{1}{\kappa_1}[y^{-1}(\theta_1 t - \kappa_1 z_1 - z_2) + \kappa_2], \tag{2.128}$$

$$\gamma = z_2 - \kappa_2(2y + \alpha - a_3), \tag{2.129}$$

$$\delta = -y^{-1}(\alpha y + z_1)(-\kappa_2 y + z_2). \tag{2.130}$$

Introduce z by

$$z_1 = \frac{y^2}{\kappa_1 q z}, \quad z_2 = \kappa_1 \kappa_2 q z (y - a_3). \tag{2.131}$$

Then the matrix $B_0(t) = (B_{ij})$ is parametrized as follows:

$$B_{11} = -q\bar{z}\left(\kappa_2 - \frac{y}{\kappa_2 \bar{z}}\right), \tag{2.132}$$

$$B_{12} = qw\bar{z}, \tag{2.133}$$

$$B_{21} = \frac{\kappa_1 q}{w(1-\kappa_1 q\bar{z})}\left(-\bar{\alpha} - \frac{\bar{y}}{\kappa_1 q\bar{z}}\right)\left(\kappa_2 - \frac{y}{\bar{z}}\right), \tag{2.134}$$

$$B_{22} = \frac{-\kappa_1 q\bar{z}}{1-\kappa_1 q\bar{z}}\left(-\bar{\alpha} - \frac{\bar{y}}{\kappa_1 q\bar{z}}\right). \tag{2.135}$$

Set further

$$b_1 = \frac{1}{\theta_1}, \quad b_2 = -\frac{\theta_1}{\kappa_1 \kappa_2 a_3}, \quad b_3 = \frac{1}{\kappa_1 q}, \quad a_4 = -\kappa_2. \tag{2.136}$$

Equation (2.120) are equivalent to

$$\frac{y\bar{y}}{a_3 a_4} = -\frac{\bar{z}(\bar{z} - b_2 t)}{\bar{z} - b_3}, \tag{2.137}$$

$$\frac{z\bar{z}}{b_3} = -\frac{y^2}{a_4(y - a_3)}, \tag{2.138}$$

$$\frac{\bar{w}}{w} = -\frac{\bar{z}}{b_3} + 1. \tag{2.139}$$

We have a constraint

$$\frac{b_1 b_2}{b_3} = q\frac{1}{a_3 a_4}. \tag{2.140}$$

q-$P(A_6)^\sharp$ is (2.137) and (2.138).

2.7 Lax form of q-$P(A_7)$

Consider a 2×2 matrix system with polynomial coefficients

$$Y(qx,t) = A(x,t)Y(x,t). \tag{2.141}$$

We express the deformation equation

$$Y(x,qt) = B(x,t)Y(x,t) \tag{2.142}$$

and can express the q-Painlevé equation of type A_7 in the form

$$A(x,qt)B(x,t) = B(qx,t)A(x,t) \tag{2.143}$$

by the compatibility of the deformation equation and the original linear q-difference equation.

We take $A(x,t)$ to be of the form

$$A(x,t) = A_0(t) + xA_1(t) + x^2 A_2, \tag{2.144}$$

$$A_2 = \begin{pmatrix} \kappa_1 & 0 \\ 0 & 0 \end{pmatrix}, \quad A_0(t) \text{ has eigenvalues } \theta_1 t,\ 0, \tag{2.145}$$

$$\det A(x,t) = \kappa_1 \kappa_2 x^3. \tag{2.146}$$

The matrix $B(x,t)$ is a rational function of the form

$$B(x,t) = \frac{1}{x}(xI + B_0(t)). \tag{2.147}$$

Define $y = y(t)$, $z_i = z_i(t)$ $(i = 1,2)$ by

$$A_{12}(y,t) = 0, \quad A_{11}(y,t) = \kappa_1 z_1, \quad A_{22}(y,t) = z_2, \tag{2.148}$$

so that

$$z_1 z_2 = \kappa_2 y^3. \tag{2.149}$$

The matrix $A(x,t)$ can be parametrized as

$$A(x,t) = \begin{pmatrix} \kappa_1((x-y)(x-\alpha) + z_1) & w(x-y) \\ \kappa_1 w^{-1}(\gamma x + \delta) & \kappa_2(x-y) + z_2 \end{pmatrix}. \tag{2.150}$$

Here

$$\alpha = \frac{1}{\kappa_1}[y^{-1}(\theta_1 t - \kappa_1 z_1 - z_2) + \kappa_2], \tag{2.151}$$

$$\gamma = z_2 - \kappa_2(2y + \alpha), \tag{2.152}$$

$$\delta = -y^{-1}(\alpha y + z_1)(\beta y + z_2). \tag{2.153}$$

Introduce z by

$$z_1 = \frac{y^2}{\kappa_1 q z}, \quad z_2 = \kappa_1 \kappa_2 q y z. \tag{2.154}$$

Then the matrix $B_0(t) = (B_{ij})$ is parametrized as follows:

$$B_{11} = -q\bar{z}\left(\kappa_2 - \frac{y}{\bar{z}}\right), \tag{2.155}$$

$$B_{12} = qw\bar{z}, \tag{2.156}$$

$$B_{21} = \frac{\kappa_1 q\bar{z}}{w(1-\kappa_1 q\bar{z})}\left(-\bar{\alpha} - \frac{\bar{y}}{\kappa_1 q\bar{z}}\right)\left(\kappa_2 - \frac{y}{\bar{z}}\right), \tag{2.157}$$

$$B_{22} = \frac{-\kappa_1 q\bar{z}}{1-\kappa_1 q\bar{z}}\left(-\bar{\alpha} - \frac{\bar{y}}{\kappa_1 q\bar{z}}\right). \tag{2.158}$$

Set further

$$b_1 = \frac{1}{\theta_1}, \quad b_2 = -\frac{\theta_1}{\kappa_1\kappa_2}, \quad b_3 = \frac{1}{\kappa_1 q}, \quad a_4 = -\kappa_2. \tag{2.159}$$

Equation (2.143) are equivalent to

$$\frac{y\bar{y}}{a_4} = \frac{b_2 t\bar{z}}{\bar{z} - b_3}, \tag{2.160}$$

$$\frac{z\bar{z}}{b_3} = -\frac{y}{a_4}, \tag{2.161}$$

$$\frac{\bar{w}}{w} = -\frac{\bar{z}}{b_3} + 1. \tag{2.162}$$

We have a constraint

$$\frac{b_1 b_2}{b_3} = q\frac{1}{a_4}. \tag{2.163}$$

q-$P(A_7)$ is (2.160) and (2.161).

2.8 Lax form of q-$P(A_7')$

Consider a 2×2 matrix system with polynomial coefficients

$$Y(qx, t) = A(x, t)Y(x, t). \tag{2.164}$$

We express the deformation equation

$$Y(x, qt) = B(x, t)Y(x, t) \tag{2.165}$$

and can express the q-Painlevé equation of type A_7' in the form

$$A(x, qt)B(x, t) = B(qx, t)A(x, t) \tag{2.166}$$

by the compatibility of the deformation equation and the original linear q-difference equation.

We take $A(x, t)$ to be of the form

$$A(x, t) = A_0(t) + xA_1(t) + x^2 A_2, \tag{2.167}$$

$$A_2 = \begin{pmatrix} \kappa_1 & 0 \\ 0 & 0 \end{pmatrix}, \quad A_0(t) \text{ has eigenvalues } \theta_1 t,\ 0, \tag{2.168}$$

$$\det A(x, t) = \kappa_1 \kappa_2 x^2. \tag{2.169}$$

The matrix $B(x, t)$ is a rational function of the form

$$B(x, t) = \frac{1}{x}(xI + B_0(t)). \tag{2.170}$$

Define $y = y(t)$, $z_i = z_i(t)$ $(i = 1, 2)$ by

$$A_{12}(y, t) = 0, \quad A_{11}(y, t) = \kappa_1 z_1, \quad A_{22}(y, t) = z_2, \tag{2.171}$$

so that

$$z_1 z_2 = \kappa_2 y^2. \tag{2.172}$$

The matrix $A(x, t)$ can be parametrized as

$$A(x, t) = \begin{pmatrix} \kappa_1((x - y)(x - \alpha) + z_1) & w(x - y) \\ \kappa_1 w^{-1}(\gamma x + \delta) & z_2 \end{pmatrix}. \tag{2.173}$$

Here

$$\alpha = \frac{1}{\kappa_1} y^{-1}(\theta_1 t - \kappa_1 z_1 - z_2), \tag{2.174}$$

$$\gamma = z_2 + \kappa_2, \tag{2.175}$$

$$\delta = -y^{-1} z_2(\alpha y + z_1). \tag{2.176}$$

Introduce z by

$$z_1 = \frac{y^2}{\kappa_1 q z}, \quad z_2 = \kappa_1 \kappa_2 q z. \tag{2.177}$$

Then the matrix $B_0(t) = (B_{ij})$ is parametrized as follows:

$$\begin{aligned} B_{11} &= qy, && (2.178)\\ B_{12} &= qw\bar{z}, && (2.179)\\ B_{21} &= \frac{-\kappa_1 qy}{w(1-\kappa_1 q\bar{z})}\left(-\bar{\alpha} - \frac{\bar{y}}{\kappa_1 q\bar{z}}\right), && (2.180)\\ B_{22} &= \frac{-\kappa_1 q\bar{z}}{1-\kappa_1 q\bar{z}}\left(-\bar{\alpha} - \frac{\bar{y}}{\kappa_1 q\bar{z}}\right). && (2.181) \end{aligned}$$

Set further

$$b_1 = \frac{1}{\theta_1}, \quad b_2 = -\frac{\theta_1}{\kappa_1\kappa_2}, \quad b_3 = \frac{1}{\kappa_1 q}, \quad b_4 = -\kappa_2. \tag{2.182}$$

Equation (2.166) are equivalent to

$$\begin{aligned} \frac{y\bar{y}}{a_4} &= -\frac{\bar{z}(\bar{z}-b_2 t)}{\bar{z}-b_3}, && (2.183)\\ \frac{z\bar{z}}{b_3} &= \frac{y^2}{a_4}, && (2.184)\\ \frac{\bar{w}}{w} &= -\frac{\bar{z}}{b_3} + 1. && (2.185) \end{aligned}$$

We have a constraint

$$\frac{b_1 b_2}{b_3} = q\frac{1}{a_4}. \tag{2.186}$$

q-$P(A_7')$ is (2.183) and (2.184).

3 Degenerations

Some replacements of the parameters for the degenerations of q-Painlevé equations were given in the paper, [6], for example. In this section, we present the replacements of the parameters of the Lax formalisms.

Replace in q-$P(A_3)$, t by εt, y by εy, z by εz, a_3 by εa_3, a_4 by $\varepsilon^{-1}a_4$ b_3 by εb_3 and b_4 by ε^{-1} and let ε tend to zero. Then we obtain q-$P(A_4)$:

$$\begin{aligned} \frac{y\bar{y}}{a_3 a_4} &= -\frac{(\bar{z}-b_1 t)(\bar{z}-b_2 t)}{\bar{z}-b_3}, && (3.1)\\ \frac{z\bar{z}}{b_3} &= -\frac{(y-a_1 t)(y-a_2 t)}{a_4(y-a_3)}. && (3.2) \end{aligned}$$

$$\frac{b_1 b_2}{b_3} = q\frac{a_1 a_2}{a_3 a_4} \tag{3.3}$$

For the sake of simplification of notation, the replacement and the succeeding limiting process will be written as follows:

$$
\begin{gathered}
t \to \varepsilon t \quad y \to \varepsilon y, \quad z \to \varepsilon z, \\
a_3 \to \varepsilon a_3, \quad a_4 \to \varepsilon^{-1} a_4 \quad b_3 \to \varepsilon b_3, \quad b_4 \to \varepsilon^{-1}.
\end{gathered}
$$

By the use of notation as above, the degeneration from the Lax form of q-$P(A_3)$ to that of q-$P(A_4)$ is given by the following scheme:

q-$P(A_3)$ to q-$P(A_4)$:

$$
\begin{gathered}
t \to \varepsilon t \quad y \to \varepsilon y, \quad z \to \varepsilon z, \\
a_3 \to \varepsilon a_3, \quad a_4 \to \varepsilon^{-1} a_4 \quad b_3 \to \varepsilon b_3, \quad b_4 \to \varepsilon^{-1}, \\
x \to \varepsilon x, \quad z_1 \to \varepsilon^2 z_1, \quad w \to \varepsilon^{-1} w, \quad \kappa_1 \to \varepsilon^{-1} \kappa_1, \quad \kappa_2 \to \varepsilon, \\
\alpha \to \varepsilon \alpha, \quad \beta \to \varepsilon^{-1} \beta, \quad \delta \to \varepsilon \delta, \\
Y(x,t) \to x^{\log_q \varepsilon} Y(x,t), \quad A(x,t) \to \varepsilon A(x,t), \\
A_0(t) \to \varepsilon A_0(t), \quad A_2 \to \varepsilon^{-1} A_2, \quad B_0(t) \to \varepsilon B_0(t), \\
B_{11} \to \varepsilon B_{11}, \quad B_{12} \to \varepsilon B_{12}, \quad B_{21} \to \varepsilon B_{21}, \quad B_{22} \to \varepsilon B_{22}.
\end{gathered}
$$

q-$P(A_4)$ to q-$P(A_5)$:

$$a_3 \to \varepsilon, \quad b_2 \to \varepsilon^{-1} b_2, \quad \theta_2 \to \varepsilon.$$

q-$P(A_4)$ to q-$P(A_5)^\sharp$:

$$
\begin{gathered}
t \to \varepsilon t, \quad a_1 \to \varepsilon^{-1} a_1, \quad a_2 \to \varepsilon, \quad b_1 \to \varepsilon b_1, \quad b_2 \to \varepsilon^{-1} b_2, \\
\theta_1 \to \varepsilon^{-1} \theta_1, \quad \theta_2 \to \varepsilon.
\end{gathered}
$$

q-$P(A_5)$ to q-$P(A_6)$:

$$t \to \varepsilon t, \quad a_1 \to \varepsilon^{-1} a_1, \quad a_2 \to \varepsilon, \quad b_1 \to \varepsilon b_1, \quad b_2 \to \varepsilon^{-1} b_2, \quad \theta_1 \to \varepsilon^{-1} \theta_1.$$

q-$P(A_5)^\sharp$ to q-$P(A_6)$:

$$a_3 \to \varepsilon, \quad b_2 \to \varepsilon^{-1} b_2.$$

q-$P(A_5)^\sharp$ to q-$P(A_6)^\sharp$:

$$a_1 \to \varepsilon, \quad b_1 \to \varepsilon b_1.$$

q-$P(A_6)$ to q-$P(A_7)$:

$$a_1 \to \varepsilon, \quad b_1 \to \varepsilon b_1.$$

q-$P(A_6)^\sharp$ to q-$P(A_7)$:

$$a_3 \to \varepsilon, \quad b_2 \to \varepsilon^{-1} b_2.$$

q-$P(A_6)^\sharp$ to q-$P(A_7')$:

$$a_3 \to \varepsilon^{-1}, \quad a_4 \to \varepsilon a_4, \quad \kappa_2 \to \varepsilon \kappa_2.$$

4 Degeneration from q-$P(A_2)$ to q-$P(A_3)$

The Lax form of q-$P(A_2)$ was given in the Sakai's paper, [8]. This Lax form yields the Lax form of q-$P(A_3)$ by a process of coalescence.

4.1 Lax form of q-$P(A_2)$

In this subsection, we illustrate the Lax form of q-$P(A_2)$ in the paper, [8].

Consider a 2×2 matrix system with polynomial coefficients

$$Y(qx,t) = A(x,t)Y(x,t). \tag{4.1}$$

We express the deformation equation

$$Y(x,qt) = B(x,t)Y(x,t) \tag{4.2}$$

and can express the q-Painlevé equation of type A_2 in the form

$$A(x,qt)B(x,t) = B(qx,t)A(x,t) \tag{4.3}$$

by the compatibility of the deformation equation and the original linear q-difference equation.

We take $A(x,t)$ to be of the form

$$A(x,t) = A_0(t) + xA_1(t) + x^2A_2(t) + x^3A_3, \tag{4.4}$$

$$A_3 = \begin{pmatrix} \kappa_1 & 0 \\ 0 & \kappa_2 \end{pmatrix}, \quad A_0(t) \text{ has eigenvalues } \theta_1 t,\ \theta_2 t, \tag{4.5}$$

$$\det A(x,t) = \kappa_1\kappa_2(x-a_1)(x-a_2)(x-a_3)(x-a_4)(x-a_5t)(x-a_6t). \tag{4.6}$$

We have

$$\kappa_1\kappa_2 a_1a_2a_3a_4a_5a_6 = \theta_1\theta_2. \tag{4.7}$$

The matrix $B(x,t)$ is a rational function of the form

$$B(x,t) = \frac{x}{(x-a_5qt)(x-a_6qt)}(xI + B_0(t)). \tag{4.8}$$

Define $\lambda = \lambda(t)$, $\mu = \mu(t)$ and $\tilde{\mu} = \tilde{\mu}(t)$ by

$$A_{12}(\lambda,t) = 0, \quad A_{11}(\lambda,t) = \kappa_1\tilde{\mu}, \quad A_{22}(\lambda,t) = \kappa_2\mu, \tag{4.9}$$

so that

$$\mu\tilde{\mu} = \kappa_1\kappa_2(\lambda-a_1)(\lambda-a_2)(\lambda-a_3)(\lambda-a_4)(\lambda-a_5t)(\lambda-a_6t). \tag{4.10}$$

The matrix $A(x,t)$ can be parametrized as

$$A(x,t) = \begin{pmatrix} \kappa_1 W(x,t) & \kappa_2 w L(x,t) \\ \kappa_1 w^{-1} X(x,t) & \kappa_2 Z(x,t) \end{pmatrix}, \tag{4.11}$$

Here

$$\begin{aligned}
L(x,t) &= x - \lambda, && (4.12)\\
Z(x,t) &= (x-\lambda)(x^2 + (\gamma+\lambda)x + \delta) + \mu, && (4.13)\\
W(x,t) &= (x-\lambda)(x^2 + (-\gamma+\lambda-\sigma_1)x + \tilde{\delta}) + \tilde{\mu}, && (4.14)\\
X(x,t) &= \frac{W(x)Z(x) - \prod_{i=1}^{6}(x-a_i)}{L(x)}, && (4.15)\\
\delta &= \frac{1}{\kappa_1-\kappa_2}\left[\kappa_1(2\lambda^2 - \sigma_1\lambda + \sigma_2 + \gamma(\gamma+\sigma_1)) - \frac{1}{\lambda}(\kappa_1\tilde{\mu} + \kappa_2\mu - \theta_1 - \theta_2)\right], && (4.16)\\
\tilde{\delta} &= \frac{1}{\kappa_1-\kappa_2}\left[-\kappa_2(2\lambda^2 - \sigma_1\lambda + \sigma_2 + \gamma(\gamma+\sigma_1)) + \frac{1}{\lambda}(\kappa_1\tilde{\mu} + \kappa_2\mu - \theta_1 - \theta_2)\right], && (4.17)\\
\tilde{\mu} &= \frac{1}{\mu}\prod_{i=1}^{6}(\lambda - a_i), \quad \sigma_1 = \sum_{i=1}^{6} a_i, \quad \sigma_2 = \sum_{i<j} a_i a_j. && (4.18)
\end{aligned}$$

If $q\kappa_1 = \kappa_2$, then Equation (4.3) are equivalent to

$$(\lambda - \underline{\nu})(\lambda - \nu) = \frac{(\lambda-a_1)(\lambda-a_2)(\lambda-a_3)(\lambda-a_4)}{(\lambda-a_5 t)(\lambda-a_6 t)}, \tag{4.19}$$

$$\left(1 - \frac{\nu}{\bar{\lambda}}\right)\left(1 - \frac{\nu}{\lambda}\right) = \frac{a_5 a_6}{q}\frac{(\nu-a_1)(\nu-a_2)(\nu-a_3)(\nu-a_4)}{(a_5 a_6 t + \theta_1/\kappa_2)(a_5 a_6 t + \theta_2/\kappa_2)}, \tag{4.20}$$

$$\begin{aligned}
a_5 a_6 t \lambda \bar{\lambda}(a_1; a_2 + a_3 + a_4 + \bar{\gamma} - \nu)((a_5+a_6)t + \gamma + \nu)& \\
+ q(a_5 a_6 t\nu + \theta_1/\kappa_2)(a_5 a_6 t\nu + \theta_2/\kappa_2) = 0.&
\end{aligned} \tag{4.21}$$

q-$P(A_2)$ is (4.19) and (4.20).

4.2 Degeneration

In this subsection, we give replacements of the parameters for the degeneration. By the use of notation in the Section 3, the degeneration from the Lax form of q-$P(A_2)$ to that of q-$P(A_3)$ is given by the following scheme:

$$\begin{gathered}
\lambda \to \varepsilon y, \quad \nu \to \varepsilon^{-1} z, \\
a_1 \to \varepsilon a_3, \quad a_2 \to \varepsilon a_4, \quad a_3 \to -\varepsilon^{-1}, \\
a_4 \to -\varepsilon^{-1} q \kappa_2^{-1} \kappa_1, \quad a_5 \to \varepsilon a_1, \quad a_6 \to \varepsilon a_2, \\
x \to \varepsilon x, \quad \mu \to \varepsilon z_2, \quad \tilde{\mu} \to \varepsilon q {\kappa_2}^{-1} \kappa_1 z_1, \quad \kappa_1 \to \varepsilon^{-1} q^{-1} \kappa_2, \quad \kappa_2 \to \varepsilon^{-1} \kappa_2, \\
\gamma \to \varepsilon^{-1} + \varepsilon \gamma_1 + O(\varepsilon^2), \\
\begin{aligned}
\gamma_1 = \frac{1}{\kappa_1 - \kappa_2} & [y^{-1}((\theta_1 + \theta_2)t - \kappa_1 z_1 - \kappa_2 z_2) \\
& - \kappa_2((a_1 + a_2)t + a_3 + a_4) + y(\kappa_1 + \kappa_2)].
\end{aligned}
\end{gathered}$$

5 Concluding remarks

By the limiting procedure, we derived Lax forms of q-Painlevé equations which are obtained from q-$P(A_3)$. The degeneration scheme from the Lax forms of q-$P(A_2)$ is also given. However Lax forms of q-$P(A_0^*)$ and q-$P(A_1)$ do not appear today, the full degeneration pattern cannot be presented. An interesting future problem remains to find the relation between the Lax pairs in the paper, [3], and our result.

Acknowledgments

The author expresses his sincere gratitude to Professor Hidetaka Sakai, who gave suggestions about ideas on this research.

References

[1] G. D. Birkhoff, The generalized Riemann problem for linear differential equations and the allied problems for linear difference and q-difference equations, *Proc. Amer. Acad. Arts Sci.* **49** (1913), 521–568.

[2] B. Grammticos, A. Ramani and V. G. Papageorgiou, Do integrable mappings have the Painlevé property?, *Phys. Rev. Lett.* **67** (1991), 1825–1828.

[3] M. Hay, J. Hietarinta, N. Joshi and F. Nijhoff, A Lax pair for a lattice modified KdV equation, reductions to q-Painlevé equations and associated Lax pairs, *J. Phys. A: Math. Theor.* **40** (2007), F61–F73.

[4] M. Jimbo and H. Sakai, A q-analog of the sixth Painlevé equation, *Lett. Math. Phys.* **38** (1996), 145–154.

[5] A. Ramani, B. Grammaticos and J. Hietarinta, Discrete versions of the Painlevé equations, *Phys. Rev. Lett.* **67** (1991), 1829–1832.

[6] A. Ramani, B. Grammaticos, T. Tamizhmani and K. M. Tamizhmani, Special function solutions of the discrete painlevé equations, *Comput. Math. Appl.* **42** (2001), 603–614.

[7] H. Sakai, A q-analog of the Garnier system, *Funkcial. Ekvac.* **48** (2005), 273–297.

[8] H. Sakai, Lax form of the q-Painlevé equation associated with the $A_2^{(1)}$ surface, *J. Phys. A: Math. Gen.* **39** (2006), 12203–12210.

[9] H. Sakai, Problem: discrete Painlevé equations and their Lax forms, *RIMS Kôkyûroku Bessatsu* **B2** (2007), 195–208.

[10] H. Sakai, Rational surfaces associated with affine root systems and geometry of the Painlevé equations, *Comm. Math. Phys.* **220** (2001), 165–229.